2. Bakgrunn (data)

2.1. Latinsk navn (Latin name)

Drymocallis rupestris (L.) Soják

2.2 Rødlistestatus (redlist satus)

(Critically endangered)

2.3 Utbredelse (spreading/place)

Drymocallis rupestris occurs in Central and South-Eastern Europe, including Scandinavia, Balkan peninsula, Northern Italy, Sardinia, Corsica, British Isles. It does not exist on other islands and near the Atlantic. Moreover, it occurs in Asia Minor, trans-Caucasia and North Africa.

2.4 Lokaliteter i Norge (locations in Norway)

2.4.1. Lokaliteter i Oslo (locations in Oslo)

1 known

Location: 1. OSLO

Individuals: 21 (Leave rosette was treated as individual)

Area: 2 m x 1.5 m (potential area 3 m x 15 m)

Environment (habitat):

SSE hill slope, too strongly shaded by old and young trees (ash – Fraxinus excelsior, elm – Ulmus sp., maple – Acer platanoides). It grows in fringe association with Geranium sanguineum (plant community) which is typical in Central Europe. Its occurence along the edges of shrub or tree stands is also typical: neighbourhood of trees and shrubs gives some shade and protection from mowing and grazing. With other species: Geranium sanguineum, Campanula trachelium, Campanula persicifolia, Filipendula vulgaris, Convallaria majalis, Polygonatum odoratum, Rosa sp., Trifolium medium, Geum urbanum, Anthriscus sylvatica, Veronica chamedrys, Festuca sp.

Condition: In 4 clusters, 3 of them with flowering plants (total: six flowering individuals)

Care: The site needs a little clearing – three closest ashes (Fraxinus) on the southern site should be cut out. Could be done gradually.

GPS-koordinates: Løvåsveien?

Date of watch: 30.06.2008

Owner:

Photos: 1673, 1675, 1676, 1678 (R. Gramsz)

Observer: R. Gramsz, J. Potocka