

Drymocallis 2014

2. Bakgrunn (data)

2.1. Latinsk navn (Latin name)

Drymocallis rupestris (L.) Soják

Leave rosette was treated as individual.

2.2 Rødlistestatus (redlist status)

(Critically endangered)

2.3 Utbredelse (spreading/place)

Drymocallis rupestris occurs in Central and South-Eastern Europe, including Scandinavia, Balkan peninsula, Northern Italy, Sardinia, Corsica, British Isles. It does not exist on other islands and near the Atlantic. Moreover, it occurs in Asia Minor, trans-Caucasia and North Africa.

2.4 Lokalteter i Norge (locations in Norway)

2.4.1. Lokalteter i Oslo (locations in Oslo)

3 known

– For maps see file [Drymocallis – Blankvann.kmz](#), [Drymocallis – Tasen.kmz](#)

[Drymocallis-Ekeberg.kmz](#)

Location: 1. OSLO – TÅSEN - 2014

Individuals: Total: 12 individuals (8 blooming + 4 vegetative).

Area: 1 m x 2 m (potential area 5 m x 40 m)

Environment (habitat):

SE hill slope, too strongly shaded by old and young trees (ash – *Fraxinus excelsior*, elm – *Ulmus glabra*, maple – *Acer platanoides*). It grows in fringe association with *Geranium sanguineum* (plant community), which is typical in Central Europe. Its occurrence along the edges of shrub or tree stands is also typical: neighbourhood of trees and shrubs gives some shade and protection from mowing and grazing. With other species: *Acer platanoides*, *Alliaria petiolata*, *Anthriscus sylvestris*, *Artemisia vulgaris*, *Campanula persicifolia*, *Campanula trachelium*, *Carex pairaei*, *Convallaria majalis*, *Dactylis glomerata*, *Filipendula vulgaris*, *Fraxinus excelsior*, *Galium boreale*, *Galium mollugo*, *Geranium sanguineum*, *Geum urbanum*, *Glechoma hederacea*, *Hylotelephium telephium*, *Festuca sp.*, *Fragaria vesca*, *Lathyrus pratensis*, *Lotus corniculatus*, *Melica nutans*, *Origanum vulgare*, *Polygonatum odoratum*, *Ranunculus acris*, *Rosa sp.*, *Rubus idaeus*, *Taraxacum officinale*, *Trifolium medium*, *Urtica dioica*, *Veronica chamedrys*, *Vicia sepium*, *Viola sp.*, *Ulmus glabra*

Condition: Only 3 clusters (1st- 1 blooming, 2nd – 6 blooming + 1 vegetative, 3rd – 1 blooming + 3 vegetative) with blooming and vegetative individuals have been found in location close to fence (phot.1.) Plants are in good condition, 20 – 50 cm. Blooming was already finished at a time of observation. We have not found any *Drymocallis* plants in location close to sand box. (Phot.2)

Care: This location is a little too much shaded by big trees and for that reason too humid in wet years.

GPS-coordinates: 59°56'42.30"N 10°44'40.01"E

[Drymocallis – Tasen.kmz](#)

Date of watch: 20.06.2014

Owner:

Photos: R. Gramsz,

Observer: R. Gramsz, J. Potocka

Phot.1. Tåsen, main location, close to fence with 3 clusters. 20.06.2014



Phot.2. Tåsen, location close to box with sand. 20.06.2014



Location: 2. OSLO – BLANKVANN – 2013

Individuals: 12 individuals (3 blooming, 9 vegetative)

Very similar as last year - in 3 clusters (2 other clusters not found)

Clusters: 1st – 6 individuals (6 vegetative), 2nd – 2 individuals (2 vegetative), 3rd – not found, 4th – not found, 5th – 4 individuals (3 blooming, 1 vegetative)

Area: 2 m x 3 m

Potential area: probably whole open deforested terrain in this place, ca. 100 m x 300 m.

Environment (habitat):

Surroundings of a Nordmarka cottage (“Stranger”) located over rocky Northern shore of Blankvann lake. Open (deforested) top and south facing hill slope. *Drymocallis rupestris* plants are located very close (10 m) to building on the area looking like already for many years running wild flower garden. On still flat area just close to hill slope, on the East side of soil hummock, partly shaded in the afternoon by rowan *Sorbus aucuparia*.

Relative of owner (both of them interested in botany) assure that this plant has not been planted – suggest it’s natural origin.

With other species: *Acer platanoides*, *Achillea millefolium*, *Agrostis capillaris*, *Alchemilla sp.*, *Anemone nemorosa*, *Antennaria dioica*, *Anthericum liliago*, *Anthoxanthum odoratum*, *Aruncus dioicus*, *Betula pubescens*, *Bergenia sp.*, *Briza media*, *Calamagrostis arundinacea*, *Campanula rotundifolia*, *Campanula trachelium*, *Convallaria majalis*, *Dryopteris filix-mas*, *Epilobium angustifolium*, *Epipactis atrorubens*, *Fragaria moschata*, *F. vesca*, *F. viridis*, *Fragaria vesca*, *Galium boreale*, *Galium verum*, *Geranium sylvaticum*, *Gymnadenia conopsea*, *Hepatica nobilis*, *Hieracium sect. hieracium*, *Hylotelephium telephium*, *Hypochoeris maculata*, *Iris sibirica*, *Lapsana communis*, *Lathyrus vernus*, *Leucantemum vulgare*, *Lilium martagon*, *Listera ovata*, *Lychnis chalconica*, *Melampyrum pratense*, *Orthilia secunda*, *Pinus sylvestris*, *Platanthera chlorantha*, *Polygala vulgaris*, *Potentilla erecta*, *Pyrola minor*, *P. rotundifolia*, *Rubus saxatilis*, *Solidago virgaurea*, *Sorbus aucuparia*, *Stachys sylvatica*, *Thymus pulegioides*, *Trifolium pratense*, *Vaccinium myrtillus*, *Vaccinium vitis-idaea*, *Valeriana officinalis*, *Vicia sp.*, *Vinca minor*, *Viola tricolor*, *Viola canina*, *Viscaria vulgaris*

Condition: Very similar as last year - *Drymocallis rupestris* is overgrown by other vegetation. Only cluster (no. 5) on top of soil hummock is flowering, the rest is very difficult to find.

Care:

GPS-coordinates: 60°01'40.8/ 010°39'57.3 [Drymocallis – Blankvann.kmz](#)

Date of watch: 21.06.2014

Owner:

Photos: R. Gramsz,

Observer: R. Gramsz, J. Potocka,

Phot.1. Blankvann, view over location of *Dryocallis rupestris*. 21.06.2014.



Phot.2. Blankvann, blooming cluster. 21.06.2014.



Phot.3. Blankvann, vegetative cluster. 21.06.2014



Location: 3. OSLO – EKEBERG – 2014

Individuals: Ca. 2000 individuals! With ca. 70% blooming.

102 clusters (tuft) in 3 concentrations (50 + 25 + 23) and 4 tufts spread some 50 m NW from main concentration.

In this much more abundant location than described above is more practical to count clusters (tufts) of rosettes. Clusters in this location are very firm, well visible and consist at average of ca. 20 rosettes each.

Area: 3 concentrations (8 x 4 m + 6 x 4 m + 5 x 3 m) grooving close together on the area not bigger than 50 x 20 m. Ca. 50m NW from main location 4 new tufts have been found.

Potential area – could be most of open area of Ekebergskrånningen Naturreservat (1 km x 50m)

Environment (habitat): SW and W rocky slop in lower (below Kongsveien) part of Ekebergskrånningen Naturreservat. Inclination of slope is between 10° to 30°. Most of area is with very shallow soil and bare rock. Cover of trees and bushes about 20% so, the location is quite open and sunny. Trees and bushes level: *Berberis vulgaris*, *Frangula alnus*, *Fraxinus excelsior*, *Pinus silvatica*, *Populus tremula*, *Prunus spinosa*, *Rhamnus cathartica*, *Rosa sp.*, *Sorbus sp.* Div., Other plants: *Achillea millefolium*, *Artemisia campestris*, *Asplenium trichomanes*, *Asplenium septentrionale* +hybrid, *Filipendula vulgaris*, *Fragaria vesca*, *Galium verum*, *Geranium sanguineum*, *Geum urbanum*, *Hylotelephium telephium*, *Hypericum perforatum*, *Inula salicyna*, *Lotus corniculatus*, *Origanum vulgare*, *Polypodium vulgare*, *Sedum album*, *Trifolium alpestre*, *Polygonatum odoratum*, *Woodsia ilvensis*, *Viscaria vulgaris*.

Condition: All plants already finish blooming at the time of observation but about 70% of leaf rosettes were blooming (or 100% of clusters have been at least with one or few blooming rosettes – usually 70% of rosettes in each cluster were blooming). Plants were usually 20 – 40 cm high, some up to 60 cm. Plants seams to be in very good condition. Only on closest to footpath concentration, 4-6 tufts have been trampled by tourists.

Care:

GPS-coordinates: 59°53'25.8"N 10°45'48.6"E

[Drymocallis-Ekeberg.kmz](#)

Date of watch: 16.06.2014; 20.06.2014

Owner:

Photos: R. Gramsz,

Observer: R. Gramsz, J. Potocka,

Phot.1. Ekeberg, the second large concentration with *Drymocallis rupestris*. 16.06.2014



Phot.2. The biggest concentration of *Drymocallis rupestris* . with some trampled tufts. 16.06.2014



Phot.3. Ekeberg, *Drymocallis rupestris*. 16.06.2014.

